

Case Report

Struma salpingi associated with Struma ovarii and steremosis – A case report

**Mansour Moghimi (MD)¹, Hamid Reza Soltani*²
Ehsan Fotouhi², Mohsen Taghavi Shavazi (MD)³**

¹*Assistant Professor, Department of Pathology, Shaheed Sadooghi University of Medical Sciences, Yazd, Iran.*

²*Medical Student, Medical Students Scientific Association of Islamic Azad University of Yazd, Yazd, Iran.*

³*General Phycisian, Medical Students Scientific Association of Islamic Azad University of Yazd, Yazd, Iran.*

Abstract

The prevalence of struma onset in uterine tube is lower than ovary's stuma. In this report a 63 year old woman was referred to main hospital in the city of Yazd, Iran in 2008. The patient referred with abdominal swelling in pre-umbilical and supra pubic region associated with losing weight and frequent constipation since 5 years ago. After the hospitalization the patient underwent the laparatomy. in pathology report the masses was struma ovarii and mature cystic teratoma associated with struma salpingi and peritoneal steremosis. Special demonstration accompanied struma salpingi in this case are struma ovarii and multiple vesicular vegetasion in choledosac, Peritoneum and bladder. This report was a very rare case of struma salpingi accompanied with struma ovarii and steremosis.

Keywords: Struma salpingi, Struma ovarii, Steremosis

* **Corresponding Author:** *Hamid Reza Soltani, E-mail: hrsgmed@yahoo.com*

Received 1 Dec 2008

Revised 18 May 2009

Accepted 28 Jun 2009